

Conversion of LKSD Styrofoam Food Service Products to a Safe Alternative

Developed by the

The Nelson Island Consortium

Analysis performed by
Zender Environmental Science and Planning Services

This discussion paper is in regards to the use of Styrofoam food service products by LKSD schools in the Nelson Island Area Villages of Chefornak, Newtown, Nightmute, Toksook Bay, and Tununak. Use of Styrofoam presents a potentially serious health threat to these communities.

The Nelson Island Consortium is dedicated to improving public health and environmental conditions in the communities and shared subsistence grounds.

Each day, Styrofoam trays, bowls, cups, and plates are used in serving lunch to LKSD students and staff. This consumption pattern results in an estimated 1.5 tons of Styrofoam discarded annually, equivalent to about 400 cubic yards of trash bags. The impact of Styrofoam waste on the environment depends on the local waste disposal method employed. It is nearly an inert product when landfilled and covered, but is toxic when incinerated, or its scattered pieces inadvertently consumed by wildlife. As with the vast majority of Alaska communities, the Nelson Island Village waste disposal sites provide only minimal protection from problematic wastes such as Styrofoam. Due to lack of cover, suitable land space, equipment, and/or operational funds, Styrofoam is ultimately burned via methods lacking emissions treatment, such as controlled open burning, burning via “burnbox”, or accidental dump fires. While allowable under State of Alaska Class 3 regulations, these methods result in the release of toxic compounds to the atmosphere, including styrene, benzene, and dioxin. Available information on smoke emissions from municipal garbage burned via these methods indicates that concentrations of these chemicals can exceed the Maximum Contaminant Levels (MCL’s) for health effect thresholds. Health effects are summarized below, with additional ecotoxicological information provided in Appendix A.

Primary contaminants emitted from the incineration of Styrofoam

Chemical	Acute Effects	Chronic Effects
Styrene	Respiratory effects, such as mucous membrane irritation, eye irritation, and gastrointestinal effects	Effects on the central nervous system (CNS), with symptoms such as headache, fatigue, weakness, depression, CNS dysfunction (reaction time, memory, visuomotor speed and accuracy, intellectual function), and hearing loss, peripheral neuropathy, minor effects on some kidney enzyme functions and on the blood.
Benzene	<i>Low-level exposure:</i> Mild reversible CNS effects, immune system depression and bone marrow toxicity leading to aplastic anemia. <i>High-level:</i> Central nervous system (CNS) effects and death	Chromosomal aberrations, cancer
Dioxin	Liver damage, weight loss, atrophy of thymus gland and immunosuppression.	Reproductive effects, including reduced fertility and birth defects

Source: Agency for Toxic Substances and Disease Registry <http://www.atsdr.cdc.gov/toxfaq.html>

Additionally, in the case of Villages where burning occurs only occasionally via dump fires, due to its flyable and breakable nature, Styrofoam litters the area surrounding the dumpsite, and can inadvertently be consumed by wildlife and fish, upon which Nelson Island residents depend for their diets.

Beyond public health concerns, due to its low specific weight, the use of Styrofoam exacts a significant, disproportionate future financial cost to Villages. Suitable land for disposal sites is very scarce, and only one to three villages each year are able to obtain funding for landfill road and facility construction statewide. Priority is for Villages of higher populations than those of Nelson Island. Styrofoam takes up approximately three to five times more volume than an equivalent weight of paper, and is non-biodegradable. Based on a YR 2005 wastestream analysis in Tununak, we estimate the school Styrofoam wastestream alone to comprise approximately 15 percent of the total community wastestream volume, as discarded. In the absence of burning the wastes, school Styrofoam would occupy approximately 8 to 10 percent of the waste volume *in situ* at Nelson Island waste disposal sites (due to natural waste compaction processes).

An issue of global responsibility for the use of Styrofoam exists as well. The production of Styrofoam consumes non-renewable petroleum products and releases toxic compounds in the nation's waterways. The manufacturing process of Styrofoam involves the use and disposal of additional chemicals, including carbon tetrachloride, polyvinyl alcohol, antimony oxide, tert-butyl hydroperoxide, and benzoquinone. When Styrofoam is produced in developing countries with relaxed environmental and worker safety regulations, workers and local communities can be exposed to these chemicals.

To evaluate the potential for such a conversion for their schools, the Nelson Island Consortium commissioned the below general cost analysis on sample food service vendor retail prices:

Styrofoam, 1000 ct.

Item	Example Product cost	Shipping Cost (\$0.37/lb)	Total Cost
Bowl, 12 oz	\$40	\$3.90	\$44
Plate, 9 "	\$50	\$8.60	\$59
Cup, 8 oz	\$30	\$3.00	\$33
Tray, 9" by 12"	\$60	\$6.00	\$66
Total Set	\$180	\$21	\$201

Alternative Technology Material, 1000 ct.

Item	Example Product cost	Shipping Cost (\$0.37/lb)	Total Cost
Bowl, 12 oz	\$45	\$8.80	\$54
Plate, 9 "	\$62	\$15.00	\$77
Cup, 8 oz	\$48	\$8.00	\$56
Tray, 9" by 12"	\$90	\$13.70	\$104
Total Set	\$245	\$45	\$291

Paper, 1000 ct.

Item	Example Product cost	Shipping Cost (\$0.37/lb)	Total Cost
Bowl, 12 oz, lightweight	\$90	\$6.40	\$96
Plate, 9 ", heavyweight	\$100	\$27.20	\$127
Cup, 8 oz, lightweight	\$58	\$5.60	\$64
Pulp Tray, 10" by 7"	\$155	\$15.50	\$171
Total Set	\$403	\$55	\$462

Our research indicates that conversion to starch-based, vegetable-oil, or other new technology material products from Styrofoam will be the least expensive alternative. Based on 185 days of food service each school year, we have calculated the following cost-to-convert to an alternative product, with an estimated 20% margin of uncertainty to account for price differences in negotiated contracts, shipping rates, and material availability and suitability:

Estimated Annual Cost To Convert From Styrofoam Bowl, Cup, Plate, And Tray To Alternative Product

Community	Number of students plus staff	Approximate number of 1,000 ct. case sets	Estimated cost to convert from Styrofoam to alternative.	Maximum estimated cost, with 20% price uncertainty for both product sets ¹
Chefornak	159	29	\$2,581	\$5,435
Newtok	119	22	\$1,958	\$4,123
Nightmute	76	14	\$1,246	\$2,624
Toksook Bay	210	38	\$3,382	\$7,121
Tununak	122	22	\$1,958	\$4,123

¹ Assumes Styrofoam set can be obtained for 20% less than estimated, and Alternative Material set must be purchased for 20% more than estimated.

Minimum operation and maintenance costs for a disposal site meeting State operational requirements to protect public health in the Nelson Island area are as follows (YR 2005 dollars):

Chefornak	Tununak	Toksook Bay	Nightmute	Newtok
\$91,134	\$91,134	\$106,700	\$75,818	\$91,134

Based on a YR 2001 feasibility study for solid waste management options in Chefornak, AK. Includes O & M of equipment, staff salary, site closure and post-care sinking fund, sinking fund for 25% match of capital costs. See Appendix B.

A YR 2001 wastestream characterization study in Chefornak determined the school contribution to the community wastestream by weight as 61% of non-residential wastes, and 7% of the total wastestream, averaged out over one calendar year¹. Assuming the school contribution is similar for each community, we provide below estimates of the school proportionate-cost share of disposal site operation and maintenance, the maximum annual cost to switch from Styrofoam to biodegradable products, and the cost difference between the two values.

Comparison of estimated Styrofoam-to-Alternative product conversion cost with school share of true costs for operation and maintenance of the local waste disposal site, annually.

Chefornak		Tununak		Toksook Bay		Nightmute		Newtok	
Maximum cost to convert	School share of waste disposal O & M	Maximum cost to convert	School share of waste disposal O & M	Maximum cost to convert	School share of waste disposal O & M	Maximum cost to convert	School share of waste disposal O & M	Maximum cost to convert	School share of waste disposal O & M
\$5,435	\$6,390	\$4,123	\$6,390	\$7,121	\$7,481	\$2,624	\$5,316	\$4,123	\$6,390
<i>Difference: \$955</i>		<i>Difference: \$2,267</i>		<i>Difference: \$360</i>		<i>Difference: \$2,692</i>		<i>Difference: \$2,267</i>	

Even using the estimated maximum cost to convert to environmentally-friendly products, our analysis indicates that a conversion is financially feasible for LKSD through mutual cooperation with the Nelson Island communities. The food service conversion is in the interest of the Nelson Island communities, and discounted disposal costs for use of the local waste site in the interest of the local schools. Our communities are willing to work with LKSD in achieving both aims.

¹ Zender Environmental Science and Planning Services, *Assessment of Solid Waste Management Situation for the Native Village of Chefornak*, funded by Central Council of Tlingit and Haida Indian Tribes of Alaska, 2001.

Appendix A:

**Summary of Ecotoxicology of Primary Contaminants Emitted from the
Open Incineration of Styrofoam**

DRAFT

Styrene vapor, a potential carcinogen, is formed from incineration of polystyrene (Styrofoam), at the low temperatures characteristic of open burning and burnboxes. Acute exposure to styrene in humans results in respiratory effects, such as mucous membrane irritation, eye irritation, and gastrointestinal effects. Based on studies of workers exposed to styrene, chronic exposure to styrene in humans results in effects on the central nervous system (CNS), with symptoms such as headache, fatigue, weakness, depression, CNS dysfunction (reaction time, memory, visuomotor speed and accuracy, intellectual function), and hearing loss, peripheral neuropathy, minor effects on some kidney enzyme functions and on the blood. No data on the health effects of breathing low-levels of styrene over long periods of time exists. And little information exists about the health effects of styrene ingestion on humans.

Used primarily as a raw material in the synthesis of styrene, benzene is a recognized carcinogen. At lower acute exposure levels to benzene above the MCL, mild CNS effects appear to be concentration dependent and rapidly reversible. But other effects include immune system depression and bone marrow toxicity leading to aplastic anemia. Acute exposure to high levels produces central nervous system (CNS) effects and death. Daily to weekly exposure to dumpsite smoke containing benzene constitutes chronic exposure to benzene. Chronic exposure to Benzene at levels above the MCL has the potential to cause chromosomal aberrations.

If benzene is released to the atmosphere, it will exist predominantly in the vapor phase. Gas-phase benzene reacts with hydroxyl radicals, resulting in the production of phenol, nitrophenols, nitrobenzene, formic acid, and peroxyacetyl nitrate. Incineration of chlorinated benzene, as occurs when Styrofoam is burned, produces dioxin. Additionally, Benzene is fairly soluble in water and is removed from the atmosphere in rain. When benzene is released to soil, it will be subject to rapid volatilization near the surface, but that which does not evaporate will be highly, to very highly, mobile in the soil and may leach to groundwater. Although most public drinking water supplies are free of benzene, exposure has been found to be very high from consumption of water sources contaminated by landfill drainage. Because Nelson Island disposal sites are unlined and proximate to drinking water sources, exposure to benzene via this pathway in addition, to the smoke inhalation pathway, is of concern.

Dioxin is one of the most toxic and environmentally stable tricyclic aromatic compounds of its structural class, and is potentially carcinogenic. Additionally, acute exposure to dioxin at levels above the MCL has been found to potentially cause liver damage, weight loss, atrophy of thymus gland and immuno-suppression. Chronic exposure to Dioxin at levels above the MCL has the potential to cause a variety of reproductive effects, including reduced fertility and birth defects.

Particulate-phase dioxin in smoke may be physically removed from air by wet and dry deposition. Due to its very low water solubility, most of the dioxin in water is expected to be associated with sediments or suspended material. Dioxin is resistant to biodegradation, and bioconcentration in aquatic organisms that consume this material has been demonstrated. Thus, contamination of subsistence resources by deposited dioxin is a potential pathway of exposure, in addition to the primary pathway of concern, smoke inhalation.

**Appendix B:
Landfill Operation and Maintenance Costs for Nelson Island Villages**

DRAFT

Estimated Operation and Maintenance Costs for Chefornak, Newtok, and Tununak⁰

Item	Unit	Cost/Unit	Quantity	Cost (\$/per year)	School contribution
Operator 0.6875 FTE ¹	hr	\$25	1430	35,750	\$2,507
Administration 0.10FTE ¹	hr	\$20	208	4,160	\$292
Equipment operation and maintenance	hr	\$23	400	9,142	\$641
Equipment replacement fund	\$93,705 for Small Track Loader, inc. shipping	\$93,705	1	3,148	\$221
Safety gear	Lump sum	\$571	1	\$571	\$40
Replacement parts (burnbox)	Lump sum	\$343	1	\$343	\$24
Generator fuel	Lump sum	\$114	1	\$114	\$8
WQ testing²	priority pollutant scan	\$1,543	1	\$1,543	\$108
Final cover/closure³	\$371,392/small site closure in bush	\$371,392	1	12,479	\$875
Post-closure care⁴	ac	\$4,000	2	2,180	\$153
Training	Lump sum	\$1,714	1	1,714	\$120
New Site Construction, match of \$964,477 for 20 yr sinking fund⁵	25%	\$241,119	1	8,102	\$568
Subtotal				\$79,246	\$5,556
Contingencies @ 15%				\$11,887	\$833
TOTAL				\$91,133	\$6,390

0 While the exact total may differ slightly, these communities share similar population sizes, and thus present similar O and M requirements.

1 Includes insurance, retirement, and administrative overhead. Equivalent to \$18 to \$20 per hr in salary.

2 Assumes testing and analysis of heavy metals and fecal coliform performed with water sampling kit (quarterly while landfill is active and once per year after closure). Priority pollutant analysis is performed once per year at an outside lab.

3 The \$371,392 is an estimate based on actual closure costs for several villages from A Guide to Closing Waste Disposal Sites in Alaska Villages, Central Council Tlingit and Haida Indian Tribes of Alaska, 2001, adjusted for ENR/CCI ratio of 7309/6396. Placed into sinking fund for planned life of landfill (20 yr) calculated at 4% interest rate, includes 40% mobilization/demobilization plus contingency rate. See note above.

4 Estimated according to actual cost average incurred for several sites. From A Guide to Closing Waste Disposal Sites in Alaska Villages, Central Council Tlingit and Haida Indian Tribes of Alaska, 2001. Placed into sinking fund for planned life of landfill (20 yr) calculated at 4% interest rate, includes 40% mobilization/demobilization plus contingency rate.

5 Placed into sinking fund of 4% interest rate.

Estimated Operation and Maintenance Costs for Toksook Bay

Item	Unit	Cost/Unit	Quantity	Cost (\$/per year)	School contribution
Operator 0.86 FTE ¹	hr	\$25	1430	\$44,688	\$3,133
Administration 0.10FTE ¹	hr	\$20	208	\$4,160	\$292
Equipment operation and maintenance	hr	\$22.85	460	\$10,513	\$737
Equipment replacement fund	\$93,705 for Small Track Loader, inc. shipping	\$93,705	1	\$3,148	\$221
Safety gear	Lump sum	\$571	1	\$571	\$40
Replacement parts (burnbox)	Lump sum	\$343	1	\$343	\$24
Generator fuel	Lump sum	\$114	1	\$114	\$8
WQ testing ²	priority pollutant scan	\$1,543	1	\$1,543	\$108
Final cover/closure ³	\$427,101/small site closure in bush	\$427,101	1	\$14,351	\$1,094
Post-closure care ⁴	ac	\$4,000	2.5	\$2,725	\$130
Training	Lump sum	\$1,714	1	\$1,714	\$120
20 yr sinking fund for capital costs of new landfill at \$1,060,924 ⁵	25% match	\$265,231	1	\$8,912	\$625
Subtotal				\$92,783	\$6,593
Contingencies @ 15%				\$13,917	\$989
TOTAL				\$106,700	\$7,582

1 Includes insurance, retirement, and administrative overhead. Equivalent to \$18 to \$20 per hr in salary.

2 Assumes testing and analysis of heavy metals and fecal coliform performed with water sampling kit (quarterly while landfill is active and once per year after closure). Priority pollutant analysis is performed once per year at an outside lab.

3 The \$427,101 is an estimate based on actual closure costs for several villages from A Guide to Closing Waste Disposal Sites in Alaska Villages, Central Council Tlingit and Haida Indian Tribes of Alaska, 2001. Placed into sinking fund for planned life of landfill (20 yr) calculated at 4% interest rate, includes 40% mobilization/demobilization plus contingency rate. See note above.

4 Estimated according to actual cost average incurred for several sites. From A Guide to Closing Waste Disposal Sites in Alaska Villages, Central Council Tlingit and Haida Indian Tribes of Alaska, 2001. Placed into sinking fund for planned life of landfill (20 yr) calculated at 4% interest rate, includes 40% mobilization/demobilization plus contingency rate.

5 YR 2001 feasibility study for solid waste management options in Cheforak, AK, plus 10% additional for larger design size of cell (note fixed costs make for economies of scale so that increase is disproportionately smaller than population ratios. Placed into sinking fund of 4% interest rate.

Estimated Operation and Maintenance Costs for Nightmute

Item	Unit	Cost/Unit	Quantity	Cost (\$/per year)	School contribution
Operator 0.52 FTE ¹	hr	\$25	1,072	\$26,813	\$1,880
Administration 0.10FTE ¹	hr	\$20	208	\$4,160	\$292
Equipment operation and maintenance	hr	\$22.85	340	\$7,771	\$545
Equipment replacement fund	\$93,705 for Small Track Loader, inc. shipping	\$93,705	1	\$3,148	\$221
Safety gear	Lump sum	\$571	1	\$571	\$40
Replacement parts (burnbox)	Lump sum	\$343	1	\$343	\$24
Generator fuel	Lump sum	\$114	1	\$114	\$8
WQ testing ²	priority pollutant scan	\$1,543	1	\$1,543	\$108
Final cover/closure ³	\$315,683/small site closure in bush	\$315,683	1	\$10,607	\$744
Post-closure care ⁴	ac	\$4,000	1.7	\$1,853	\$130
Training	Lump sum	\$1,500	1	\$1,714	\$120
20 yr sinking fund for capital costs of new landfill at \$868,029 ⁵	25% match	217,007	1	\$7,292	\$511
Subtotal				\$65,929	\$4,623
Contingencies @ 15%				\$9,889	\$693
TOTAL				\$75,818	\$5,316

1 Includes insurance, retirement, and administrative overhead. Equivalent to \$18 to \$20 per hr in salary.

2 Assumes testing and analysis of heavy metals and fecal coliform performed with water sampling kit (quarterly while landfill is active and once per year after closure). Priority pollutant analysis is performed once per year at an outside lab.

3 The \$315,683 is an estimate based on actual closure costs for several villages from A Guide to Closing Waste Disposal Sites in Alaska Villages, Central Council Tlingit and Haida Indian Tribes of Alaska, 2001. Placed into sinking fund for planned life of landfill (20 yr) calculated at 4% interest rate, includes 40% mobilization/demobilization plus contingency rate. See note above.

4 Estimated according to actual cost average incurred for several sites. From A Guide to Closing Waste Disposal Sites in Alaska Villages, Central Council Tlingit and Haida Indian Tribes of Alaska, 2001. Placed into sinking fund for planned life of landfill (20 yr) calculated at 4% interest rate, includes 40% mobilization/demobilization plus contingency rate.

5 Based on YR 2001 feasibility study for solid waste management options in Chefornak, AK, minus 10% costs for smaller design size of cell (note fixed costs make for economies of scale so that decrease is disproportionately smaller than population ratios. Placed into sinking fund of 4% interest rate.